

OR-25

REARRANGEMENT OF FURYL-TETHERED OXIMES: SYNTHESIS OF FUNCTIONALIZED PYRROLES


A. S. Makarov, A. A. Fadeev, Y.A. Vasev, M.G. Uchuskin

Perm State University, 15 Bukireva St., Perm, 614990, Russia.

E-mail: antony.s.makarov@psu.ru

Abstract. Polyarylated pyrroles have been extensively utilized as building-blocks in material science.¹ High demand for functionalized pyrrole substrates motivated us to develop a general synthetic protocol toward such valuable compounds based on furan dearomatization approach.²

We report here a novel dearomatizative rearrangement of furyl-tethered oxime esters that provides 2,4-di(het)arylpyrroles possessing highly reactive acylvinyl fragment.



Scheme 1. Synthesis of functionalized pyrroles

Scope and limitations of the described protocol as well as further synthetic applications of the obtained products will be discussed.

References

1. Loudet A. BODIPY Dyes and Their Derivatives: Syntheses and Spectroscopic Properties / A. Loudet, K. Burgess // Chemical Reviews. – 2007. – Vol. 107, Iss. 11. – P. 4891–4932.
2. Trushkov I. V. Furan's gambit: Electrophile-attack-triggered sacrifice of furan rings for the intramolecular construction of azaheterocycles / I. V. Trushkov, M. G. Uchuskin, A. V. Butin // European Journal of Organic Chemistry. – 2015. – P. 2999–3016.

This work was supported by the Russian Foundation for Basic Research, project # 20-33-70079.